

# CO<sub>2</sub> and associated parameters variability in the Southern Ocean: CARIOCA and ARGO drifters

J. Boutin (1), L. Merivat (1), C. Henocq (1)  
 Laboratoire d'Océanographie et du Climat par Expérimentation et Analyse Numérique / Institut Pierre Simon Laplace / Université Pierre et Marie Curie, France

In the frame of the EU-FP6 CARBOOCEAN project, 19 months of CARIOCA data have been acquired by four CARIOCA buoys deployed in the Southern Atlantic Ocean in 2005, 2006 and 2007.  
 In this poster, we analyse the variability of pCO<sub>2</sub> with respect to 1) Takahashi (2002) climatology, 2) the location of the drifters relative to the subantarctic front and 3) the biological activity.

## CARIOCA MEASUREMENTS IN THE SOUTHERN ATLANTIC OCEAN SINCE 2005

CARIOCA drifters measure each hour, at 2 meters depth, ocean CO<sub>2</sub> partial pressure, pCO<sub>2</sub>, sea surface temperature, SST, salinity, SSS, fluorescence, and in the atmosphere surface wind speed, atmospheric pressure and air temperature.




Fig 1: 2005, 2006 and 2007 CARIOCA measurements. From top to bottom: pCO<sub>2</sub>, SSS and SST.

## COMPARISON WITH TAKAHASHI (2002) CLIMATOLOGY

Takahashi et al. (2002) derived a pCO<sub>2</sub> climatology referred to 1995 year, from ship measurements. We extract pCO<sub>2</sub> values from this climatology corresponding to location and months of CARIOCA measurements.

Table: Difference between pCO<sub>2</sub> from Takahashi (2002) referred to 1995 year and pCO<sub>2</sub> measured by CARIOCA drifters. Standard deviation of the differences are also indicated.

	pCO <sub>2</sub> Taka - pCO <sub>2</sub> Carioca (µatm)
2005-North East	-24 ± 12
2006-2007 South	-31 ± 10
2006-2007 North-East	-9 ± 23
2007 North-Mid Atlantic	-26 ± 8

pCO<sub>2</sub> is always undersaturated wrt atmospheric pCO<sub>2</sub> (370 ± 5 µatm over that period). It exhibits a large variability; pCO<sub>2</sub> as low as 280 µatm are observed.

On average over 3 CARIOCA tracks, pCO<sub>2</sub> from Takahashi climatology is 24 to 31 µatm lower than pCO<sub>2</sub> measured by CARIOCA drifters. This difference is significant (well above the standard deviation of the differences): this is consistent with an increase of pCO<sub>2</sub> in surface waters that follows the atmospheric trend (20 µatm between 1995 and 2006; N.B. Takahashi (2002) did not correct data south of 50°S for atmospheric trend so that the difference may be larger than 20 µatm).

The difference with the 2006-2007 North-East buoy is much weaker and much more variable because of strong biological activity (see below)

## TAKAHASHI(2002) CLIMATOLOGY ALONG CARIOCA TRACKS




Fig 2: Takahashi (2002) pCO<sub>2</sub> climatology along CARIOCA tracks. Color scale has been shifted by 20 µatm wrt CARIOCA pCO<sub>2</sub> figure in order to take into account oceanic trend.

Once the atmospheric trend is taken into account, the east-west gradient observed on the Southern buoy with lower pCO<sub>2</sub> in Summer-Fall between 50W and 30W is similar to Takahashi (2002) climatology.

On another hand, the strong difference between 2006 north-east and south buoys between 10E and 25E is not observed on the climatology

## LOCATION OF CARIOCA MEASUREMENTS RELATIVE TO SUBANTARCTIC FRONT

2006 CARIOCA remains south of the Subantarctic zone, in polar zone, as indicated by its low SSS (SSS < 34) and SST. Climatological fronts (Orsi et al., 1995; Belkin and Gordon 1996) were found too approximative to precisely locate CARIOCA with respect to Subantarctic front, SAF. We detect SAF from Microwave Optimally Interpolated (OI) Sea Surface Temperature (SST) maps (available on [www.ssmi.com/SST/microwave\\_o\\_i\\_sst\\_browse.html](http://www.ssmi.com/SST/microwave_o_i_sst_browse.html)) as the maximum SST gradient within 5°C and 9°C (as proposed by Moore et al. (1999) and Burls and Reason (2006)).




Fig 3: South and North-East CARIOCA trajectories superimposed on AMSR SST map (29/11/06: CARIOCA north-east deployment)




Fig 4: SSS recorded by CARIOCA 2006 South and North-East and by ARGO profilers (color points) superimposed on SAF detected from maximum SST gradient (black).

## SAF SST GRADIENT

Both CARIOCA SSS, ARGO SSS (Salinity gradient around 34.1, indicative of SAF) and SST gradient criteria indicate that:  
 -2006 South buoy was south of SAF (Fig4)  
 -2006-2007 North East buoy was deployed on the SAF (Fig4)  
 -2005 and 2007 buoys were deployed in the SAZ (Fig1)

## BIOLOGICAL ACTIVITY ALONG CARIOCA TRAJECTORIES

The low pCO<sub>2</sub> observed by the 2006 north-east buoy (Fig 1) are associated with very large fluorescence values (Fig 5). They correspond to a chlorophyll filament observed onto the SAF at the time of the buoy deployment (Fig 6, left) and to large chlorophyll contents at the edge of an eddy when the buoy drifts northwards (Fig 6, right).






Fig 5: CARIOCA measurements of fluorescence (daily averages).

Fig 6: CARIOCA trajectory superimposed on (left) SEAWIFS chlorophyll image just after the buoy deployment, (right) MODIS chlorophyll image when the buoy was in the north east of the image



## SUMMARY

A CARIOCA drifter records pCO<sub>2</sub> for more than one year in the Polar Zone of the Atlantic Ocean; the mean air-sea CO<sub>2</sub> flux derived with the Wanninkhof (1992) and the Ho et al. (2006) k-U relationship is 1.2 mol m<sup>-2</sup> yr<sup>-1</sup> and 1. mol m<sup>-2</sup> yr<sup>-1</sup> respectively; the minimum pCO<sub>2</sub> is observed in Summer-Fall between 50W and 30W and is associated with strong fluorescence values.

The 3 other buoys drift north of the SAF. The minimum pCO<sub>2</sub> (down to 280 µatm) is observed in November 06 when the 2006 North-East buoy was on the SAF and in January-February 2007: during both periods, the buoy follows thin chlorophyll filaments. This strengthens the small scale variability of pCO<sub>2</sub> in the open ocean associated with strong chlorophyll variability linked to ocean dynamics.

CARIOCA pCO<sub>2</sub> is on average larger than Takahashi(2002) climatology by a value close to the atmospheric trend between 1995 and 2006, confirming that the surface ocean, even at high latitudes, follows the atmospheric trend (Takahashi, 2006).

Acknowledgements: CARIOCA preparations were supervised by L. Beaumont, T. Danguy and V. Dutreuil from DT INSU. We thank N. Martin for computation support and L. Barbero for very constructive remarks. We are indebted to M. Hoppema, O. Klatt, L. Beaumont, E. Isla, C. Bock, P. Sangiardi, M. Calzas, N. Metz and to Polarstern and Marion Dufresne/IPEV crews for CARIOCA deployments.

References:  
 Burls, N.J., and C.J.C. Reason, Sea surface temperature fronts in the midlatitude South Atlantic revealed by using microwave satellite data, *J. Geophys. Res.*, 111, C08001, doi:10.1029/2005JC003133, 2006.  
 Moore, J.K., M.R. Abbott, and J.G. Richman, Location and dynamics of the Antarctic Polar Front from satellite sea surface temperature data, *Journal of Geophysical Research*, 104, C2, 3059-3073, 1999.  
 Takahashi, T., S.C. Sutherland, C. Sweeney, A. Poisson, N. Metz, B. Tilbrook, N. Bates, R. Wanninkhof, R.A. Feely, C. Sabine, J. Olafsson, and Y. Nojiri, Global sea-air CO<sub>2</sub> flux based on climatological surface ocean pCO<sub>2</sub> and seasonal biological and temperature effects, *Deep Sea Research Part II*, 49 (9-10), 1601-1622, 2002.  
 Takahashi, T., and al., Climatological mean surface ocean pCO<sub>2</sub> and net air-sea CO<sub>2</sub> flux over the global oceans: progress since 2002, 2006, available at [www.whoi.edu/cms/files/mzawoysky/2006/8/takahashi\\_ocb\\_July06\\_13065.pdf](http://www.whoi.edu/cms/files/mzawoysky/2006/8/takahashi_ocb_July06_13065.pdf).